

Addendum to “Groups of Ribbon Knots”

by Ka Yi Ng

The purpose of this document is to clarify the inductive step described in the proof of Theorem 3.2 in [2]. In the second last sentence of the proof, it says, ‘it follows from the inductive proof that $\mathcal{R}_n$ is of index two.’ The question of how this assertion is verified was first raised by Dror Bar Natan and his student Ofer Ron [1]. To avoid any confusions that may arise in the future, the author of the paper [2] would like to fill in the details to show that $\mathcal{R}_n$ is of index two. Readers of this document are assumed to have familiarity with [2]. For your reference, Theorem 3.2 is stated below.

Theorem 3.2 $\mathcal{R}_n$ forms a subgroup of the free abelian group $\mathcal{V}_n$ of index two. So its rank is the same as the rank of $\mathcal{V}_n$ and is the number of linearly independent primitive rational invariants of order $\leq n$.

To show that $\mathcal{R}_n$ is of index two, we shall establish the following two lemmas.

Lemma A: Let $D_1, \dots, D_r$ be all the distinct non-split chord diagrams of order $n \geq 2$. Let $\lambda_1, \dots, \lambda_r$ be r arbitrary integers. Then for $n > 2$, there exists a ribbon knot K_n whose additive invariants of order $< n$ are trivial and $V(K_n) = \sum_{i=1}^r \lambda_i V(D_i)$ for all additive invariants V of order n . When $n = 2$, there exists a ribbon knot K_2 with $V(K_2) = 2\lambda_1 V(D_1)$ for all additive invariants V of order 2.

Lemma B: Given a knot K whose additive invariants of order $< n$ are trivial, there exist n -chord diagrams $D_1, \dots, D_r$ and integers $\lambda_1, \dots, \lambda_r$ such that $V(K) = \sum_{i=1}^r \lambda_i V(D_i)$ for all additive invariants V of order n .

Lemma A is an extended version of Lemma 3.1 in [2] and can be proved using the same arguments as given in [2]. Lemma B is proved in [3]. We shall now make use of these two lemmas to prove Lemma C.

Lemma C: Let X be the right-handed trefoil. Let K be an arbitrary knot and let $n \geq 2$ be an integer. Define s to be 1 if the Arf invariant of the knot K is non-zero and 0 otherwise. Then one can construct a ribbon knot K_n so that $V(K \# sX \# K_n) = 0$ for all additive invariants V of order $\leq n$.

Proof of Lemma C: For $n = 2$, the knot can always be represented by an integral multiple λ of the non-split 2-chord diagram D so that $V(K) = \lambda V(D)$ for all additive invariants of order 2. Write λ as $2p - s$ where p is an integer. By Lemma A, we can find a ribbon knot K_2 with $V(K_2) = -2pV(D)$ for all additive invariants of order 2. The right-handed trefoil knot has $V(X) = V(D)$ and thus, the case $n = 2$ is proved.

Suppose a ribbon knot K_n is constructed so that $V(K \# sX \# K_n) = 0$ for all additive invariants of order $\leq n$. Lemma B tells us that there exist integers $\lambda_1, \dots, \lambda_r$

and $(n + 1)$ -chord diagrams $D_1, \dots, D_r$ so that $V(K \# sX \# K_n) = \sum_{i=1}^r \lambda_i V(D_i)$ for all additive invariants of order $n + 1$. Apply Lemma A to construct a ribbon knot R whose additive invariants of order $\leq n$ are trivial and $V(R) = -\sum_{i=1}^r \lambda_i V(D_i)$ for all additive invariants of order $n + 1$. Then $V(K \# sX \# K_n \# R) = 0$ for all additive invariants of order $\leq n + 1$.

Lemma C implies that $V(K \# sX \# K_n) = 0$ for all rational-valued invariants of order $\leq n$. According to the proof of Theorem 3.2 in [2], $\mathcal{R}_n$ is a subgroup of $\mathcal{V}_n$. Thus, as an element in $\mathcal{V}_n$, we have $[K]_n \in -[sX]_n + \mathcal{R}_n$. This completes the proof that $\mathcal{R}_n$ is of index two.

References

- [1] Dror Bar Natan and Ofen Ron, *A correction to "Groups of Ribbon Knots" by Ka Yi Ng*, arXiv:math.GT/0309282 17 Sep 2003.
- [2] K.Y.Ng, *Groups of Ribbon Knots*, Topology **37**, No.2, 441-458 (1998).
- [3] K.Y.Ng and T.Stanford, *On Gusarovs groups of knots*, Proc. Camb Phil Soc. **126**, No. 1, 63-76 (1999).

Ka Yi Ng, e-mail: Kayi.Ng@WallStreetSystems.com